arXiv:1510.00358v3 [astro-ph.HE] 7 Jul 2016

Observation of the new emission line at ∼3.5 keV in X-ray spectra of galaxies and galaxy clusters

D. A. Iakubovskyi[1,2]

[1]Discovery Center, Niels Bohr Institute, Blegdamsvej 17, Copenhagen, Denmark
[2]Bogolyubov Institute of Theoretical Physics, Metrologichna Str. 14-b, 03680, Kyiv, Ukraine
iakubovskyi@nbi.ku.dk

The detection of an unidentified emission line in X-ray spectra of cosmic objects would be a 'smoking gun' signature for particle physics beyond the Standard Model. More than a decade of its extensive searches results in several narrow faint emission lines reported at 3.5, 8.7, 9.4 and 10.1 keV. The most promising of them is the emission line at ∼3.5 keV reported in spectra of several nearby galaxies and galaxy clusters. Here I summarize its up-to-date status, overview its possible interpretations, including an intriguing connection with radiatively decaying dark matter, and outline future directions for its studies.

Key words: X-rays: general, dark matter, line: identification.

INTRODUCTION

The origin of *dark matter* – the major (yet of unknown origin) gravitating substance in the Universe [1–25] – still has to be revealed. If dark matter is made of elementary particles, the corresponding particle should be massive (to form over-densities in process of gravitational collapse), long-lived (to be stable for at least the age of the Universe) and neutral with respect to strong and electromagnetic interactions (to be sufficiently 'dark'). The only known massive, long-lived and neutral particles are the usual (left-handed) neutrinos, but they are too light to form small dark matter haloes [26, 27]. As a result, the hypothesis of dark matter particle implies an extension of the Standard Model of particle physics. Dozens of the Standard Model extensions proposed so far to contain a valid dark matter particle candidate. However, as Fig. 1 of [28] demonstrates, the masses of dark matter particle candidates and their interaction strengths with Standard Model particles cover a huge region of parameter space. This results in a large variety of observational methods developed to search for dark matter particles.

The specific example considered in this review is *radiatively decaying dark matter*. If a dark matter particle interacts with electrically charged particles, it may [1] possess a radiative decay channel. If a non-relativistc dark matter particle decays to a photon and another particle, slight ($v/c \lesssim 5 \times 10^{-3}$) Doppler broadening due to non-zero velocities of dark matter particles in halos would cause a narrow *dark matter decay line*. Such a decay line possesses several specific features allowing to robustly distinct it from emission lines of astrophysical origin (see e.g. [29, 30]) or from instrumental line-like features:

- its position in energy is solely determined by the mass of dark matter particle and the redshift of dark matter halo (i.e. if one neglects the mass of the other decay product, the line position is $\frac{m_{\rm DM}c^2}{2(1+z)}$), having different scaling with halo redshift z compared with instrumental line-like features;
- its intensity is proportional to *dark matter column density* $\mathcal{S}_{\rm DM} = \int \rho_{\rm DM} d\ell$; due to different 3D distributions of dark and visible matter, comparison of the new line intensity within the given object and among different objects would allow to choose between its decaying dark matter and astrophysical origins;
- it is broadened with the characteristic velocity of dark matter different from that of visible matter.

The above-mentioned characteristics allow to *directly detect the radiatively decaying dark matter relying on astrophysical measurements*. This motivates the extensive search for new lines in X-ray spectra of cosmic objects proposed about 15 years ago [31–33], see Table 1. An example is the analysis of the line candidate at ∼2.5 keV initially reported by [34] in X-ray spectrum of the Willman 1 dwarf spheroidal at 2.5σ level. Further non-observation of this line candidate in central part and outskirts of Andromeda galaxy, Fornax and Sculptor dwarf spheroidal galaxies [35] excludes the decaying dark matter origin of the ∼2.5 keV signal at high significance level (above 14σ). This result is further strengthened by the authors of [36] who reanalyzed the same observations of Willman 1 as [34] (and did not find the ∼2.5 keV line) and the authors of [37] who analysed another dwarf spheroidal, Segue 1. Finally, the authors of [38] ruled out the dark matter origin of the ∼2.5 keV feature by looking at Willman 1 with better statistics. The probable origin of the ∼2.5 keV line, according to [35], is purely instrumental, being the result of under-modelling of the time-variable soft proton background (see e.g. [39]) in some observations combined with an apparent dip at ∼2.5 keV in the effective area of existing X-ray instruments.

Observational evidence for the line at ∼3.5 keV

The new emission line at ∼3.5 keV is reported by two different groups [70, 71] in February 2014.

In [70], the authors combine X-ray emission from the sample of nearby galaxy clusters observed by the European Photon and Imaging Camera (EPIC) [72, 73] on-board the *XMM-Newton* X-ray cosmic observatory [74] with the largest number of counts ($> 10^5$ counts for redshifts $z < 0.1$ and $> 10^4$ counts for redshifts $0.1 < z < 0.4$). The stacking is made in the cluster's rest frame. As a result, the emission from instrumental lines is smeared out, while cosmic lines appear more prominent. This method allows [70] to detect 28 emission lines of astrophysical origin in 2-10 keV band, much more than in individual galaxy clusters, see e.g. [75]. Apart of them, [70] identify the new line located at 3.57±0.02 keV in *XMM-Newton*/MOS [72] cameras and at 3.51±0.03 keV in *XMM-Newton*/PN [73] camera at the level $\gtrsim 10$ larger than predicted by two complexes of nearby astrophysical emission lines located at 3.51 keV (K XVIII)

[1] The widely-known examples where this is *not* the case are dark matter particles are the *lightest* particles holding a new quantum number *conserved* by the Standard Model interactions, such as R-parity for super-symmetric models, Kaluza-Klein number for extra dimensions, etc.. In this case, dark matter decays are strictly forbidden by the special structure of the theory, and the main astrophysical effect for dark matter particles is their *annihilation* with their antiparticles.

Ref.	Object	Instrument	Cleaned exposure, ks
[40]	Diffuse X-ray background	HEAO-1, *XMM-Newton*/EPIC	224, 1450
[41]	Coma, Virgo	*XMM-Newton*/EPIC	20, 40
[42]	Large Magellanic Cloud	*XMM-Newton*/EPIC	20
[43]	Milky Way	*Chandra*/ACIS-S3	Not specified
[44]	M31 (central $5'$)	*XMM-Newton*/EPIC	35
[45]	Abell 520	*Chandra*/ACIS-S3	67
[46]	Milky Way, Ursa Minor	*XMM-Newton*/EPIC	547, 7
[47]	Milky Way	*Chandra*/ACIS	1500
[48]	1E 0657-56 ("Bullet cluster")	*Chandra*/ACIS-I	450
[49]	Milky Way	X-ray micro-calorimeter	0.1
[50]	Milky Way	INTEGRAL/SPI	5500
[51]	M31 (central $5 - 13'$)	*XMM-Newton*/EPIC	130
[52]	Milky Way	INTEGRAL/SPI	12200
[53]	Ursa Minor	*Suzaku*/XIS	70
[54]	Draco	*Chandra*/ACIS-S	32
[34]	Willman 1	*Chandra*/ACIS-I	100
[35]	M31, Fornax, Sculptor	*XMM-Newton*/EPIC , *Chandra*/ACIS	400, 50, 162
[36]	Willman 1	*Chandra*/ACIS-I	100
[37]	Segue 1	Swift/XRT	5
[55]	M33	*XMM-Newton*/EPIC	20-30
[56]	M31 ($12 - 28'$ off-centre)	*Chandra*/ACIS-I	53
[38]	Willman 1	*XMM-Newton*/EPIC	60
[57]	Ursa Minor, Draco	*Suzaku*/XIS	200, 200
[58]	Stacked galaxies	*XMM-Newton*/EPIC	8500
[59]	M31	*Chandra*/ACIS-I	404
[60]	Stacked dSphs	*XMM-Newton*/EPIC	410
[61]	Stacked galaxies	*XMM-Newton*/EPIC, *Chandra*/ACIS-I	14600, 15000
[62]	Perseus	*Suzaku*/XIS	520
[63, 64]	Milky Way	Fermi/GBM	4600
[65]	Milky Way	*Suzaku*/XIS	31500
[66]	Draco	*XMM-Newton*/EPIC	87
[67]	1E 0657-56 ("Bullet cluster")	NuSTAR	266
[68]	Draco	*XMM-Newton*/EPIC	1660

Table 1: Summary of searches for dark matter decay line in X-ray observations conducted so far. This Table is an update of Table 1 in [69].

and 3.62 keV (Ar XVII). The new line is also detected at $> 3\sigma$ local significance in several different sub-samples of their combined *XMM-Newton*/EPIC cluster dataset, see e.g. Fig. 1, and in *Chandra*/ACIS spectrum of Perseus cluster, see Table 2 for details.



Fig. 1: The combined MOS spectrum of Perseus cluster scaled to 3-4 keV energy range. On top of their best-fit model, the series of the single-bin residuals corresponding to the extra emission line at 3.57 keV is shown in red. (Adapted from Figure 7 in [70]).

The authors of [71] detect the new line at 3.53±0.03 keV in the central part of Andromeda galaxy (see Fig. 2), and in the outskirts of Perseus cluster, see Table 2. [71] exclude the central part of Perseus cluster (analysed in [70]) because of its rather complex structure in X-rays, so the two datasets used in [70, 71] are totally independent enhancing the statistical significance for the new line. Another important result of [71] is the radial dependence of the new line flux in Perseus that appears more consistent with decaying dark matter profile than with astrophysical emission.

The encouraging results of [70, 71] have stimulated several groups to look on other dark matter-dominated objects. The following searches report the presence of the line at ∼3.5 keV, see Table 3:

1. The identification of the line at ∼3.5 keV from the region of Galactic Centre [78–81]. Although it is unclear whether the detected line has astrophysical origin (see the next Sec. for detailed discussion), its explanation in terms of decaying dark matter is consistent with the previous new line detections, see [80, 82] for details.

2. The detection of the new line in *Suzaku*/XIS observations of Perseus, Coma and Ophiuchus galaxy clusters [83]. While subsequent study of *Suzaku*/XIS spectra by [62] does not reveal the new line at ∼3.5 keV in the central part of Perseus cluster, another recent study by [84] does; however, its apparent discrepancy with the negative result of [62] is still unclear and has to be resolved further.

3. The detection of the new line at 3.52±0.08 keV observed in X-ray spectra of 8 individual nearby galaxy clusters including Perseus and Coma [85].



Fig. 2: The same as in the previous Figure 1 but for the combined spectrum of Andromeda galaxy. (Adapted from Figure 1 in [71]).

In summary, positive detections of the new line listed in Table 2 and Table 3 support the hypothesis of decaying dark matter implying radiatively decaying dark matter lifetime $\tau_{\rm DM} \simeq (6-8) \times 10^{27}$ s [80, 85, 89].

On the contrary, the following studies do not detect the ∼3.5 keV line putting the upper bounds on its flux:

1. The central part of the Virgo cluster observed by *Chandra*/ACIS [70], *Suzaku*/XIS[83] and *XMM-Newton*/EPIC [85], as well as other 10 galaxy clusters from [85].

2. Combined spectrum from dwarf spheroidal galaxies [60].

3. Outskirts of galaxies [58, 61, 71].

4. Combined blank-sky observations [65, 71].

5. Prolonged *XMM-Newton*/EPIC observations of Draco dwarf spheroidal galaxy [68, 89]; although the authors of [89] report a line-like excess at 3.54±0.06 keV with $\Delta\chi^2 = 5.3$ in PN camera, see Table 2, this finding is not supported by an independent analysis of [68] and is not accompanied with similar excess in Draco spectra seen by MOS camera [68, 89].

Ref.	Object	Redshift	Instrument	Exposure, Ms	Line position, keV	Line flux, 10^{-6} ph/s/cm^2
[70]	Full stacked sample	0.009-0.354	MOS	6	3.57±0.02	4.0±0.8
[70]	Full stacked sample	0.009-0.354	PN	2	3.51±0.03	$3.9^{+0.6}_{-1.0}$
[70]	Coma+Centaurus+Ophiuchus	0.009-0.028	MOS	0.5	3.57[a]	$15.9^{+3.4}_{-3.8}$
[70]	Coma+Centaurus+Ophiuchus	0.009-0.028	PN	0.2	3.57[a]	< 9.5 (90%)
[70]	Perseus (< 12')	0.016	MOS	0.3	3.57[a]	$52.0^{+24.1}_{-15.2}$
[70]	Perseus (< 12')	0.016	PN	0.05	3.57[a]	< 17.7 (90%)
[70]	Perseus (1-12')	0.016	MOS	0.3	3.57[a]	$21.4^{+7.0}_{-6.3}$
[70]	Perseus (1-12')	0.016	PN	0.05	3.57[a]	< 16.1 (90%)
[70]	Rest of the clusters	0.012-0.354	MOS	4.9	3.57[a]	$2.1^{+0.4}_{-0.5}$
[70]	Rest of the clusters	0.012-0.354	PN	1.8	3.57[a]	$2.0^{+0.3}_{-0.5}$
[70]	Perseus (> 1')	0.016	ACIS-S	0.9	3.56±0.02	$10.2^{+3.7}_{-3.5}$
[70]	Perseus (< 9')	0.016	ACIS-I	0.5	3.56[a]	$18.6^{+7.8}_{-8.0}$
[70]	Virgo (< 500")	0.003-0.004	ACIS-I	0.5	3.56[a]	< 9.1 (90%)
[71]	M31 (< 14')	-0.001[b]	MOS	0.5	3.53±0.03	$4.9^{+1.6}_{-1.3}$
[71]	M31 (10-80')	-0.001[b]	MOS	0.7	3.50-3.56	< 1.8 (2σ)
[71]	Perseus (23-102')	0.0179[b]	MOS	0.3	3.50±0.04	7.0±2.6
[71]	Perseus (23-102')	0.0179[b]	PN	0.2	3.46±0.04	9.2±3.1
[71]	Perseus, 1st bin (23-37')	0.0179[b]	MOS	0.2	3.50[a]	13.8±3.3
[71]	Perseus, 2nd bin (42-54')	0.0179[b]	MOS	0.1	3.50[a]	8.3±3.4
[71]	Perseus, 3rd bin (68-102')	0.0179[b]	MOS	0.03	3.50[a]	4.6±4.6
[71]	Blank-sky	—	MOS	7.8	3.45-3.58	< 0.7 (2σ)

Table 2: Properties of the ∼3.5 keV line reported by [70, 71]. For their analysis, the authors of [70, 71] use different X-ray datasets observed by MOS [73] and PN [72] cameras on-board *XMM-Newton* observatory [74] and ACIS instrument [76] on-board *Chandra* observatory [77]. All error bars are at 1σ (68%) level.
[a] The line position is fixed at given value.
[b] The redshift is fixed at NASA Extragalactic Database (NED) value.

Ref.	Object	Redshift	Instrument	Exposure, Ms	Line position, keV	Line flux, 10^{-6} ph/s/cm^2
[78]	Galactic centre (2.5-12')	0.0	ACIS-I	0.8	3.51	$\simeq 10^a$
[79]	Galactic centre (0.3-15')	0.0	MOS	0.7	3.51	45 ± 4^a
[79]	Galactic centre (0.3-15')	0.0	PN	0.5	3.51	39 ± 7^a
[79]	M31	0.0	MOS	0.5	3.53±0.07	2.1±1.5^c
[80]	Galactic centre (< 14')	0.0	MOS	0.7	3.539±0.011	29±5
[83]	Perseus core (< 6')	0.0179^b	XIS	0.74	$3.510^{+0.023}_{-0.008}$	$32.5^{+3.7}_{-4.3}$
[83]	Perseus confined (6-12.7')	0.0179^b	XIS	0.74	$3.510^{+0.023}_{-0.008}$	$32.5^{+3.7}_{-4.3}$
[83]	Coma (< 12.7')	0.0231^b	XIS	0.164	$\simeq 3.45^d$	$\simeq 30^d$
[83]	Ophiuchus (< 12.7')	0.0280^b	XIS	0.083	$\simeq 3.45^d$	$\simeq 40^d$
[83]	Virgo (< 12.7')	0.0036^b	XIS	0.09	3.55^a	< 6.5 (2σ)
[85]	Abell 85 (< 14')	0.0551^b	MOS	0.20	$3.44^{+0.06}_{-0.05}$	$6.3^{+3.9}_{-3.6}$
[85]	Abell 2199 (< 14')	0.0302^b	MOS	0.13	$3.41^{+0.04}_{-0.04}$	$10.1^{+5.1}_{-4.8}$
[85]	Abell 496 (< 14')	0.0329^b	MOS	0.13	$3.55^{+0.06}_{-0.09}$	$7.5^{+6.1}_{-4.4}$
[85]	Abell 496 (< 14')	0.0329^b	PN	0.08	$3.45^{+0.04}_{-0.03}$	$16.8^{+5.9}_{-6.4}$
[85]	Abell 3266 (< 14')	0.0589^b	PN	0.06	$3.53^{+0.04}_{-0.06}$	$8.7^{+5.1}_{-4.5}$
[85]	Abell S805 (< 14')	0.0139^b	PN	0.01	$3.63^{+0.05}_{-0.06}$	$17.1^{+9.3}_{-7.4}$
[85]	Coma (< 14')	0.0231^b	MOS	0.17	$3.49^{+0.04}_{-0.05}$	$23.7^{+10.7}_{-9.0}$
[85]	Abell 2319 (< 14')	0.0557^b	MOS	0.08	$3.59^{+0.05}_{-0.06}$	$18.6^{+10.7}_{-7.4}$
[85]	Perseus (< 14')	0.0179^b	MOS	0.16	$3.58^{+0.05}_{-0.08}$	$25.2^{+12.5}_{-12.6}$
[85]	Virgo^e (< 14')	0.0036^b	PN	0.06	—	< 9.3
[89]	Draco (< 14')	0.0	PN	0.65	$3.54^{+0.06}_{-0.05}$	$1.65^{+0.67}_{-0.70}$
[84]	Perseus (< 8.3')	0.0179^b	XIS	1.67	3.54±0.01	$27.9^{+3.5}_{-3.5}$
[84]	Perseus (< 2')	0.0179^b	XIS	1.67	3.51±0.02	$9.3^{+2.6}_{-2.7}$
[84]	Perseus (2'-4.5')	0.0179^b	XIS	1.67	3.55±0.02	$16.7^{+2.9}_{-3.0}$
[84]	Perseus (4.5'-8.3')	0.0179^b	XIS	1.67	3.58±0.02	$16.1^{+3.2}_{-3.4}$
[90]	Stacked clusters	0.01-0.45	XIS	8.1	3.54^f	$1.0^{+0.5}_{-0.5}$

Table 3: Properties of ∼3.5 keV line searched after February 2014 in different X-ray datasets observed by MOS [73] and PN [72] cameras on-board *XMM-Newton* observatory [74], ACIS [76] instrument on-board *Chandra* observatory [77] and XIS instrument [91] on-board *Suzaku* observatory [92]. All error bars are at 1σ (68%) level.
[a] Best-fit line flux at fixed position 3.51 keV coinciding with the brightest K XVIII line.
[b] Redshift was fixed at NASA Extragalactic Database (NED) value.
[c] The line is detected at < 90% confidence level. Such a low flux (compared with [71]) is because of non-physically enhanced level of continuum at 3-4 keV band used in [79], see [93] for details.
[d] Parameters estimated from Fig. 3 of [83].
[e] Given an example of the new line non-detection, see Table II of [85] for more details.
[f] Line position is fixed at the best-fit energy detected in *Suzaku* observations of the Perseus cluster by [84].



Fig. 3: The flux of the ∼3.5 keV line from the Galactic Centre, the Perseus cluster outskirts, the Andromeda galaxy, and the 'blank sky' dataset [71] as a function of dark matter projected mass. Diagonal lines show the expected behaviour of decaying dark matter signal for a given dark matter particle lifetime. The vertical sizes of the boxes are $\pm 1\sigma$ statistical error on the line's flux – or the 2σ upper bound for the blank-sky dataset. The blue shaded regions show a particular Navarro-Frenk-White [86, 87] profile of the Milky Way [88], its horizontal size indicates uncertainties in galactic disk modelling. The lifetime $\tau_{\rm DM} \sim (6-8) \times 10^{27}$ s is consistent with all datasets. New results from a prolonged Draco *XMM-Newton*/EPIC observation [68, 89] give controversial results: while [68] reports an exclusion of dark matter hypothesis at 99% level, the results of [89] claim that the values of $\tau_{\rm DM} \simeq (7-9) \times 10^{27}$ sec are still consistent with *all* existing observations. (Adapted from Figure 2 in [80]).



Fig. 4: The same as in previous Fig. 3 but over-plotted are the ranges for the $> 2\sigma$ detections in MOS (green) and PN (magenta) cameras, see [85]. (Adapted from Figure 2 in [85]).

6. Combined dataset of 33 galaxy clusters observed by *Chandra*/ACIS [94].

At the moment, it is unclear whether these negative searches rule out the decaying dark matter hypothesis of the new line. While the bounds obtained in [60] are mildly consistent with the decaying dark matter origin of the detections in [70, 71], the results of [61] formally exclude the decaying dark matter hypothesis of the ∼3.5 keV line imposing the very strict 3σ bound, $\tau_{\rm DM} > 1.8 \times 10^{28}$ s. Taking into account systematic effects in spectra (e.g. causing significant negative residuals) obtained by [61] and the apparent uncertainty in their dark matter distributions [95] would result in much weaker bound, see e.g. $\tau_{\rm DM} \gtrsim 3.5 \times 10^{27}$ s [96] using the stacked dataset of nearby galaxies of [58] with comparable exposure, still consistent with existing detections. The uncertainty in dark matter distributions also helps to reconcile the results of the other negative searches [56, 59, 65] with ∼3.5 keV line detections using the decaying dark matter paradigm. There is also no clarity with the new prolonged (∼ 1.4 Ms) *XMM-Newton*/EPIC observation of Draco dwarf spheroidal galaxy – the object having both well-measured dark matter distribution [97] and proven low X-ray background [54, 60, 82, 98]. While [68] reports an exclusion of dark matter hypothesis at 99% level having 2σ upper bound on radiative dark matter decay lifetime $\tau_{\rm DM} > 2.7 \times 10^{28}$ s, the results of [89] suggest $\tau_{\rm DM} \simeq (7-9) \times 10^{27}$ s, the value still compatible with all existing observations.

"Standard" explanations of the line at ∼3.5 keV

There are three possible "standard" explanations of the new line detections at ∼3.5 keV:

1. statistical fluctuations;
2. general-type systematic effects;
3. astrophysical emission line.

With recent increase of positive detections reported by [85], it is very hard to explain *all* of the detections with pure statistical fluctuations. Nevertheless, statistical fluctuations may be responsible for new line detections or non-detections in *some* individual objects, as well as for variations of the detected line position up to ∼110 eV [85], see Fig. 5 – the effect that should be properly taken into account when searching for the new line (unlike [60, 61, 83]).

The systematic origin of the line is carefully investigated because of the previous study of the line-like residual at ∼2.5 keV in the Willman 1 dwarf



Fig. 5: The position of new line detected in [85] (in the frame of emitting galaxy cluster) as a function of cluster redshift. The red and black dashed lines show the expected behaviour in case of purely systematic and cosmic line origins (assuming the line position 3.52 keV in the detector frame expected from [71, 80]), respectively. (Adapted from Figure 3 in [85]).

spheroidal, see Sec. for details. However, the explanation of the ∼3.5 keV line with general-type systematics suggested in [79] is unlikely. For example, its position (in the frame of emitting object) remains remarkably constant with redshift [70, 71, 85], see Fig. 5, which cannot be explained by simple systematics. The line is also independently detected by five detectors on-board three cosmic missions, see Table 2 and Table 3. Finally, similar feature of systematic origin should be detected in the blank-sky dataset [71], and should have different radial behaviour in the outskirts of Perseus cluster [71, 84].

On the other hand, the explanation of the new line with the K XVIII line complex at ∼3.5 keV suggested by [79] (see also an extensive discussion in [70, 93, 99, 100]) is still possible, at least for Galactic Centre region and galaxy clusters, contrary to initial claims of [70, 71]. The reason is that the emission flux from the K XVIII line complex at ∼3.51 keV suggested by [79] is highly uncertain due to large uncertainties of the Potassium abundance, see e.g. [101, 102] for a potential[2] level of uncertainty. Moreover, unlike other possible emission lines of astrophysical origin near ∼3.5 keV (such as Cl XVII lines at 3.51 keV found largely sub-dominant in Galactic Centre region [79] and in galaxy clusters [99]), K XVIII line complex does not have stronger counterparts at other energies and can hardly be excluded by measurements of other lines, the strongest of them is the K XIX line complex at 3.71 keV of comparable strength [103]. The same is true about the charge exchange of S XVI ions recently suggested by [104].

An alternative approach is to study the *line morphology*. At the moment, two different methods have been used. The first method [70, 71] is to split the region covered by astrophysical sources onto several independent subregions, large enough to detect the line at in each of them, and to model their spectra separately looking for a line-like excess in each of them. As a result, [70] show that the ∼3.5 keV line in Perseus cluster is somewhat more concentrated compared to decayed dark matter distributed according to Navarro-Frenk-White [86, 87] profile. By studying the ∼3.5 keV line emission from Perseus cluster outskirts, [71] obtain that such distribution is better consistent with radiatively decaying dark matter distributed according to the well-established Navarro-Frenk-White profile than with astrophysical continuum emission distributed according to the isothermal β-model of [105]. The recent detailed study [84] confirms this result and expands it to the central region of Perseus cluster.

The second method to study the line morphology [81] deals with spatial distribution of the 'line plus continuum' X-ray emission in Perseus cluster and Galactic Centre region with further eliminating continuum component by either assuming it spatially smooth or cross-correlating the 'line plus continuum' images in several energy bands (including those dominated by astrophysical line emission). By using the second method, the authors of [81] show that adding decaying dark matter distribution from a *smooth* dark matter profile (Navarro-Frenk-White, Einasto, Burkert) does not improve the fit quality in both objects, and demonstrate that distribution of the events in 3.45-3.6 keV bands correlates with that in the energy bands of strong astrophysical emission, rather than with that in line-free energy bands. Based on these findings, Ref. [81] claims the exclusion of decaying dark matter origin of 3.5 keV in Galactic Centre and Perseus cluster.

To ultimately check the astrophysical origin of the ∼3.5 keV line, new observations with high-resolution imaging [3] spectrometers such as Soft X-ray Spectrometer (SXS) [108] on-board the recently launched *Hitomi*[4] (former *Astro-H*) mission [111], *Micro-X* sounding rocket experiment [112] and the X-ray Inte-

[2] The results of [102] indicate an order of magnitude overabundance of Potassium in solar corona compared to solar photosphere. Based on this result, [102] suggested that the Potassium abundance in hot plasma in galaxies and galaxy clusters may have also been enhanced compared to the solar photospheric values. However, because at the moment there is no established mechanism that could effectively provide such an enhancement, the results of [102] only indicate the *potential* level of uncertainty, similar to the measurements in [101].

[3] *Grating* spectrometers such as *Chandra*/HETGS [106] have excellent spectral resolution for *point* sources; however, for extended ($\gtrsim$1 arcmin) sources their spectral resolution usually degrades to that for existing imaging spectrometers, see e.g. [107].

[4] Although *Hitomi* is now broken apart, it had observed Perseus cluster before the break-up [109, 110].



Fig. 6: Line emissivities (in arbitrary units) broadened with energy resolution of Soft X-ray Spectrometer (SXS) on-board *Hitomi* (former *Astro-H*), $\sigma_{SXS} = 5$ eV, as functions of energy for three-component model of [79] of Galactic Centre. The relative S, Ar, Cl and K abundances are set to $1/3 : 1 : 1 : 3$, according to Sec. 2.2 of [79]. Thin dashed line shows the total line emissivity. (Adapted from Figure 2 in [103]).

gral Field Unit (X-IFU) [113, 114] on-board planned *Athena* mission [115, 116]. If the position of the new line incidentally coincides with that of K XVIII line complex, more detailed study of the ratios of the Potassium line emissivities will be essential to finally check the astrophysical origin of the new line, see Fig. 6 for details.

Other extra line candidates in X-ray range

Although the line at ∼3.5 keV receives the largest attention of the community, there are three other line candidates in X-rays which origin is also not established:

1. According to [117], intensity of the *Fe XXVI Ly-γ line* at 8.7 keV observed in *Suzaku*/XIS spectrum of the Milky Way centre [118] cannot be explained by standard ionization and recombination processes, and dark matter decay may be a possible explanation of this excess.

2. According to Sec. 1.4 of [119], two faint extra line-like excesses at 9.4 and 10.1 keV are detected in the combined *Suzaku*/XIS spectrum of Galactic Bulge region. Notably, positions of these excesses do not coincide with any bright[5] astrophysical or instrumental line and their intensities can be explained in frames of decaying dark matter origin, see right Fig. 8 of [119].

[5]The newest available atomic database AtomDB v.3.0.2 [120] contains several faint Ni XXVI - Ni XXVIII emission lines at 10.02-10.11 keV.

Possible implications for new physics

If none of "conventional" explanations discussed in the previous Sec. were valid, the existence of the new line at ∼3.55 keV will be an indication of a new physics beyond the Standard Model.

Historically, the first model discussed in connection with ∼3.5 keV detection is the neutrino minimal extension of the Standard Model with three right-handed (sterile) neutrinos (the νMSM) [121–123]. In this model, the lightest sterile neutrino with mass in keV range forms the bulk of dark matter while two heavier sterile neutrinos are responsible for two other established phenomena beyond the Standard Model – neutrino oscillations and generation of asymmetry between baryons and anti-baryons in early Universe. Sterile neutrinos decay possess the 2-body radiative channel $N \to \gamma + \nu$, so the observation of ∼3.5 keV decay line would imply the existence of light sterile neutrino dark matter particles with mass ∼7.1 keV. The simplest production scenario of sterile neutrino dark matter – via non-resonant oscillations of usual (active) neutrinos in the early Universe [31–33, 124–126] – is already excluded by the combination of X-ray measurements [51], measurements of Lyman-α forest [127–132] and the phase-space bound from dwarf spheroidal galaxies [26, 133–136]. The realistic scenario of dark matter production within the νMSM now involves resonant oscillations of active neutrinos in hot primeval plasma with significant lepton asymmetry generated by decays of heavier sterile neutrinos [137–141]. The parameters of observed ∼3.5 keV line are consistent with νMSM predictions, see Fig. 7 for details. Because the interaction of sterile neutrino dark matter with Standard Model particles is orders of magnitude weaker than that of ordinary neutrinos, its prospects for direct detection in a particle physics experiment are very far from the existing experimental technique, see [142–146]. To confirm the νMSM, a search for heavier sterile neutrinos in GeV range is needed, handled by e.g. planned Search for Hidden Particles (*SHiP*) experiment [147, 148] and Future electron-positron e^+e^- Circular Collider (*FCC-ee*) [149].

However, the confirmation of decaying dark matter origin of the new line does not imply the existence of νMSM sterile neutrinos as there are plenty of other alternatives which can potentially explain the ∼3.55 keV line, see e.g. [80, 96, 146] and the references therein. Differences among these models can be further probed by:

- changes in the new line morphology because of non-negligible initial dark matter velocities, see e.g. [150, 151];



Fig. 7: Constraints on sterile neutrino dark matter within the νMSM model [121–123]. In every point in the white region sterile neutrinos constitute 100% of dark matter and their properties agree with the existing bounds. The blue point corresponds to the observed line from Andromeda galaxy, while the error bars indicate statistical errors (thick) and uncertainty in dark matter distribution at the central part of Andromeda galaxy (thin). (Adapted from Figure 4 in [71]).

- other astrophysical and cosmological tests, see e.g. [63, 133, 139, 152–165];
- search for "smoking gun" signatures in future dedicated particle physics experiments, such as *SHiP* [147, 148] and *FCC-ee* [149] experiments.

Recently proposed alternatives to radiatively decaying dark matter include: decay of excited dark matter states [166–174], annihilating dark matter [175–178], dark matter decaying into axion-like particles with further conversion to photons in magnetic field [179–183]. These models predict *substantial* difference in ∼3.5 keV line morphology compared to the radiatively decaying dark matter. For example, the spatial distributions of the new line in these models should be more concentrated towards the centres of dark matter-dominated objects compared to radiatively decaying dark matter, e.g. due to larger dark matter density (for excited and annihilating dark matter) or larger magnetic fields (for magnetic field conversion of axion-like particles). Further non-observation of the ∼3.5 keV line in outskirts of dark matter-dominated objects would argue in favour of these models.

Conclusion and future directions

The origin of the new emission line at ∼3.5 keV reported by [70, 71, 80, 83, 85] remains unexplained. The observed properties of the new line are consistent with radiatively decaying dark matter and other interesting scenarios (such as, exciting dark matter, annihilating dark matter and dark matter decaying into axion-like particles further converted in cosmic magnetic fields) motivated by various particle physics extensions of the Standard Model. In case of radiatively decaying dark matter, further detections would lead to direct detection of new physics. Specially dedicated observations using existing X-ray missions (such as *XMM-Newton*, *Chandra*, *Suzaku*) still allow such detections although one should take detailed care on various systematic effects that could mimic or hide the new line.

The alternative is to use new better instruments. The basic requirements for such instruments – higher *grasp* (the product of field-of-view and effective area) and better *spectral resolution* – have first formulated in [49]. Both the soft X-ray Spectrometer [108] on-board the new X-ray mission *Hitomi* (former *Astro-H*) [111, 184] and the planned *Micro-X* sounding rocket experiment [112] meet only second requirement having the energy resolution by an order of magnitude better (∼ 5 eV) than existing imaging spectrometers. Before being broken apart, *Hitomi* has already observed Perseus cluster [109]. It is expected [70] that such an observation would allow *Hitomi* to precisely determine the new line position in brightest objects with prolonged observations and to detect the K XIX emission line complex at ∼3.71 keV. Another possible option is to resolve the intrinsic width of the new line because of its Doppler broadening in galaxies and galaxy clusters [70, 185]. As a result, *Hitomi*/SXS is able to check whether the new line comes from new physics or from (anomalously enhanced) astrophysical emission. The same is expected from the *Micro-X* rocket-based microcalorimeter (to be launched in 2017) which will observe the central region of our Galaxy. Another possibility is to use the planned *eROSITA* instrument on-board *Spektrum-Röntgen-Gamma* mission [186] and the planned *LOFT* mission [187] which high grasp and moderate energy resolution would allow to detect the new line at much smaller intensities [69, 188]. Finally, an "ultimate" imaging spectrometer proposed in e.g. [189] (an example is the X-ray Integral Field Unit (X-IFU) [113, 114] on-board the planned *Athena* mission [115, 116]) would reveal the detailed morphology structure of the ∼3.5 keV line [190].

Acknowledgement

This work was supported by a research grant from VILLUM FONDEN. The author also acknowledges partial support from the Swiss National Science Foundation grant SCOPE IZ7370-152581, the Program of Cosmic Research of the National Academy of Sciences of Ukraine, the State Fund for Fundamental Research of Ukraine and the State Programme of Implementation of Grid Technology in Ukraine dur-

ing early stages of this work.

REFERENCES

[1] Zwicky, F. Die Rotverschiebung von extragalaktischen Nebeln. *Helvetica Physica Acta* **6**, 110–127 (1933).

[2] Sarazin, C. L. X-ray emission from clusters of galaxies. *Reviews of Modern Physics* **58**, 1–115 (1986).

[3] Evrard, A. E., Metzler, C. A. & Navarro, J. F. Mass Estimates of X-Ray Clusters. ApJ **469**, 494 (1996). astro-ph/9510058.

[4] Einasto, J. & Einasto, M. Dark Matter in Groups and Clusters of Galaxies. In Valtonen, M. J. & Flynn, C. (eds.) *IAU Colloq. 174: Small Galaxy Groups*, vol. 209 of *Astronomical Society of the Pacific Conference Series*, 360 (2000). astro-ph/9909437.

[5] Bergström, L. Non-baryonic dark matter: observational evidence and detection methods. *Reports on Progress in Physics* **63**, 793–841 (2000). hep-ph/0002126.

[6] Corbelli, E. Dark matter and visible baryons in M33. MNRAS **342**, 199–207 (2003). astro-ph/0302318.

[7] Refregier, A. Weak Gravitational Lensing by Large-Scale Structure. ARA&A **41**, 645–668 (2003). astro-ph/0307212.

[8] Dekel, A. *et al.* Lost and found dark matter in elliptical galaxies. Nature **437**, 707–710 (2005). astro-ph/0501622.

[9] Massey, R. *et al.* Dark matter maps reveal cosmic scaffolding. Nature **445**, 286–290 (2007). astro-ph/0701594.

[10] Gilmore, G. *et al.* The Observed Properties of Dark Matter on Small Spatial Scales. ApJ **663**, 948–959 (2007). astro-ph/0703308.

[11] Noordermeer, E., van der Hulst, J. M., Sancisi, R., Swaters, R. S. & van Albada, T. S. The mass distribution in early-type disc galaxies: declining rotation curves and correlations with optical properties. MNRAS **376**, 1513–1546 (2007). astro-ph/0701731.

[12] Fu, L. *et al.* Very weak lensing in the CFHTLS wide: cosmology from cosmic shear in the linear regime. A&A **479**, 9–25 (2008). 0712.0884.

[13] Coccato, L. *et al.* Kinematic properties of early-type galaxy haloes using planetary nebulae. MNRAS **394**, 1249–1283 (2009). 0811.3203.

[14] Einasto, J. Dark Matter. *ArXiv e-prints* (2009). 0901.0632.

[15] Rozo, E. *et al.* Cosmological Constraints from the Sloan Digital Sky Survey maxBCG Cluster Catalog. ApJ **708**, 645–660 (2010). 0902.3702.

[16] Reid, B. A. *et al.* Cosmological constraints from the clustering of the Sloan Digital Sky Survey DR7 luminous red galaxies. MNRAS **404**, 60–85 (2010). 0907.1659.

[17] Chemin, L., Carignan, C. & Foster, T. H I Kinematics and Dynamics of Messier 31. ApJ **705**, 1395–1415 (2009). 0909.3846.

[18] Corbelli, E., Lorenzoni, S., Walterbos, R., Braun, R. & Thilker, D. A wide-field H I mosaic of Messier 31. II. The disk warp, rotation, and the dark matter halo. A&A **511**, A89 (2010). 0912.4133.

[19] Massey, R., Kitching, T. & Richard, J. The dark matter of gravitational lensing. *Reports on Progress in Physics* **73**, 086901 (2010). 1001.1739.

[20] Tinker, J. L. *et al.* Cosmological Constraints from Galaxy Clustering and the Mass-to-number Ratio of Galaxy Clusters. ApJ **745**, 16 (2012). 1104.1635.

[21] Roos, M. Astrophysical and Cosmological Probes of Dark Matter. *Journal of Modern Physics* **3**, 1152–1171 (2012). 1208.3662.

[22] Frenk, C. S. & White, S. D. M. Dark matter and cosmic structure. *Annalen der Physik* **524**, 507–534 (2012). 1210.0544.

[23] Hinshaw, G. *et al.* Nine-year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Parameter Results. ApJS **208**, 19 (2013). 1212.5226.

[24] Walker, M. *Dark Matter in the Galactic Dwarf Spheroidal Satellites*, 1039 (2013).

[25] Planck Collaboration *et al.* Planck 2015 results. XIII. Cosmological parameters. *ArXiv e-prints* (2015). 1502.01589.

[26] Tremaine, S. & Gunn, J. E. Dynamical role of light neutral leptons in cosmology. *Physical Review Letters* **42**, 407–410 (1979).

[27] White, S. D. M., Frenk, C. S. & Davis, M. Clustering in a neutrino-dominated universe. ApJ **274**, L1–L5 (1983).

[28] Gardner, S. & Fuller, G. M. Dark matter studies entrain nuclear physics. *Progress in Particle and Nuclear Physics* **71**, 167–184 (2013). 1303.4758.

[29] Dere, K. P., Landi, E., Mason, H. E., Monsignori Fossi, B. C. & Young, P. R. CHIANTI - an atomic database for emission lines. A&AS **125**, 149–173 (1997).

[30] Smith, R. K., Brickhouse, N. S., Liedahl, D. A. & Raymond, J. C. Collisional Plasma Models with APEC/APED: Emission-Line Diagnostics of Hydrogen-like and Helium-like Ions. ApJ **556**, L91–L95 (2001). astro-ph/0106478.

[31] Dolgov, A. D. & Hansen, S. H. Massive sterile neutrinos as warm dark matter. *Astroparticle Physics* **16**, 339–344 (2002). hep-ph/0009083.

[32] Abazajian, K., Fuller, G. M. & Patel, M.

Sterile neutrino hot, warm, and cold dark matter. Phys. Rev. D **64**, 023501 (2001). astro-ph/0101524.

[33] Abazajian, K., Fuller, G. M. & Tucker, W. H. Direct Detection of Warm Dark Matter in the X-Ray. ApJ **562**, 593–604 (2001). astro-ph/0106002.

[34] Loewenstein, M. & Kusenko, A. Dark Matter Search Using Chandra Observations of Willman 1 and a Spectral Feature Consistent with a Decay Line of a 5 keV Sterile Neutrino. ApJ **714**, 652–662 (2010). 0912.0552.

[35] Boyarsky, A. *et al.* Searching for dark matter in X-rays: how to check the dark matter origin of a spectral feature. MNRAS **407**, 1188–1202 (2010). 1001.0644.

[36] Mirabal, N. & Nieto, D. Willman 1 in X-rays: Can you tell a dwarf galaxy from a globular cluster? *ArXiv e-prints* (2010). 1003.3745.

[37] Mirabal, N. Swift observation of Segue 1: constraints on sterile neutrino parameters in the darkest galaxy. MNRAS **409**, L128–L131 (2010). 1010.4706.

[38] Loewenstein, M. & Kusenko, A. Dark Matter Search Using XMM-Newton Observations of Willman 1. ApJ **751**, 82 (2012). 1203.5229.

[39] Kuntz, K. D. & Snowden, S. L. The EPIC-MOS particle-induced background spectra. A&A **478**, 575–596 (2008).

[40] Boyarsky, A., Neronov, A., Ruchayskiy, O. & Shaposhnikov, M. Constraints on sterile neutrinos as dark matter candidates from the diffuse X-ray background. MNRAS **370**, 213–218 (2006). astro-ph/0512509.

[41] Boyarsky, A., Neronov, A., Ruchayskiy, O. & Shaposhnikov, M. Restrictions on parameters of sterile neutrino dark matter from observations of galaxy clusters. Phys. Rev. D **74**, 103506 (2006). astro-ph/0603368.

[42] Boyarsky, A., Neronov, A., Ruchayskiy, O., Shaposhnikov, M. & Tkachev, I. Strategy for Searching for a Dark Matter Sterile Neutrino. *Physical Review Letters* **97**, 261302–+ (2006). astro-ph/0603660.

[43] Riemer-Sørensen, S., Hansen, S. H. & Pedersen, K. Sterile Neutrinos in the Milky Way: Observational Constraints. ApJ **644**, L33–L36 (2006). astro-ph/0603661.

[44] Watson, C. R., Beacom, J. F., Yüksel, H. & Walker, T. P. Direct x-ray constraints on sterile neutrino warm dark matter. Phys. Rev. D **74**, 033009–+ (2006). astro-ph/0605424.

[45] Riemer-Sørensen, S., Pedersen, K., Hansen, S. H. & Dahle, H. Probing the nature of dark matter with cosmic x rays: Constraints from "dark blobs" and grating spectra of galaxy clusters. Phys. Rev. D **76**, 043524–+ (2007). astro-ph/0610034.

[46] Boyarsky, A., Nevalainen, J. & Ruchayskiy, O. Constraints on the parameters of radiatively decaying dark matter from the dark matter halos of the Milky Way and Ursa Minor. A&A **471**, 51–57 (2007). astro-ph/0610961.

[47] Abazajian, K. N., Markevitch, M., Koushiappas, S. M. & Hickox, R. C. Limits on the radiative decay of sterile neutrino dark matter from the unresolved cosmic and soft x-ray backgrounds. Phys. Rev. D **75**, 063511 (2007). astro-ph/0611144.

[48] Boyarsky, A., Ruchayskiy, O. & Markevitch, M. Constraints on Parameters of Radiatively Decaying Dark Matter from the Galaxy Cluster 1E 0657-56. ApJ **673**, 752–757 (2008). astro-ph/0611168.

[49] Boyarsky, A., den Herder, J., Neronov, A. & Ruchayskiy, O. Search for the light dark matter with an X-ray spectrometer. *Astroparticle Physics* **28**, 303–311 (2007). astro-ph/0612219.

[50] Yüksel, H., Beacom, J. F. & Watson, C. R. Strong Upper Limits on Sterile Neutrino Warm Dark Matter. *Physical Review Letters* **101**, 121301–+ (2008). 0706.4084.

[51] Boyarsky, A., Iakubovskyi, D., Ruchayskiy, O. & Savchenko, V. Constraints on decaying dark matter from XMM-Newton observations of M31. MNRAS **387**, 1361–1373 (2008). 0709.2301.

[52] Boyarsky, A., Malyshev, D., Neronov, A. & Ruchayskiy, O. Constraining dark matter properties with SPI. MNRAS **387**, 1345–1360 (2008). 0710.4922.

[53] Loewenstein, M., Kusenko, A. & Biermann, P. L. New Limits on Sterile Neutrinos from Suzaku Observations of the Ursa Minor Dwarf Spheroidal Galaxy. ApJ **700**, 426–435 (2009). 0812.2710.

[54] Riemer-Sørensen, S. & Hansen, S. H. Decaying dark matter in the Draco dwarf galaxy. A&A **500**, L37–L40 (2009).

[55] Borriello, E., Paolillo, M., Miele, G., Longo, G. & Owen, R. Constraints on sterile neutrino dark matter from XMM-Newton observations of M33. MNRAS **425**, 1628–1632 (2012). 1109.5943.

[56] Watson, C. R., Li, Z. & Polley, N. K. Constraining sterile neutrino warm dark matter with Chandra observations of the Andromeda galaxy. *JCAP* **3**, 18 (2012). 1111.4217.

[57] Kusenko, A., Loewenstein, M. & Yanagida, T. T. Moduli dark matter and the search for its decay line using Suzaku x-ray telescope. Phys. Rev. D **87**, 043508 (2013). 1209.6403.

[58] Iakubovskyi, D. *Constraining properties of dark matter particles using astrophysical data.* Ph.D. thesis, Instituut-Lorentz for Theoretical Physics (2013).

[59] Horiuchi, S. *et al.* Sterile neutrino dark matter

bounds from galaxies of the Local Group. *Phys. Rev. D* **89**, 025017 (2014). 1311.0282.

[60] Malyshev, D., Neronov, A. & Eckert, D. Constraints on 3.55 keV line emission from stacked observations of dwarf spheroidal galaxies. Phys. Rev. D **90**, 103506 (2014). 1408.3531.

[61] Anderson, M. E., Churazov, E. & Bregman, J. N. Non-detection of X-ray emission from sterile neutrinos in stacked galaxy spectra. MNRAS **452**, 3905–3923 (2015). 1408.4115.

[62] Tamura, T., Iizuka, R., Maeda, Y., Mitsuda, K. & Yamasaki, N. Y. An X-ray spectroscopic search for dark matter in the Perseus cluster with Suzaku. PASJ **67**, 23 (2015). 1412.1869.

[63] Horiuchi, S., Ng, K. C. Y., Gaskins, J. M., Smith, M. & Preece, R. Improved limits on sterile neutrino dark matter from full-sky observations by the Fermi-GBM. *ArXiv e-prints* (2015). 1502.03399.

[64] Ng, K. C. Y., Horiuchi, S., Gaskins, J. M., Smith, M. & Preece, R. Improved limits on sterile neutrino dark matter using full-sky Fermi Gamma-ray Burst Monitor data. Phys. Rev. D **92**, 043503 (2015). 1504.04027.

[65] Sekiya, N., Yamasaki, N. Y. & Mitsuda, K. A search for a keV signature of radiatively decaying dark matter with Suzaku XIS observations of the X-ray diffuse background. PASJ (2015). 1504.02826.

[66] Sonbas, E. *et al.* X-Ray Sources in the Dwarf Spheroidal Galaxy Draco. ApJ **821**, 54 (2016). 1505.00216.

[67] Riemer-Sørensen, S. *et al.* Dark Matter Line Emission Constraints from NuSTAR Observations of the Bullet Cluster. ApJ **810**, 48 (2015). 1507.01378.

[68] Jeltema, T. & Profumo, S. Deep XMM observations of Draco rule out at the 99 per cent confidence level a dark matter decay origin for the 3.5 keV line. MNRAS **458**, 3592–3596 (2016). 1512.01239.

[69] Neronov, A., Boyarsky, A., Iakubovskyi, D. & Ruchayskiy, O. Potential of the Large Observatory for X-ray Timing telescope for the search for dark matter. Phys. Rev. D **90**, 123532 (2014). 1312.5178.

[70] Bulbul, E. *et al.* Detection of an Unidentified Emission Line in the Stacked X-Ray Spectrum of Galaxy Clusters. ApJ **789**, 13 (2014). 1402.2301.

[71] Boyarsky, A., Ruchayskiy, O., Iakubovskyi, D. & Franse, J. Unidentified Line in X-Ray Spectra of the Andromeda Galaxy and Perseus Galaxy Cluster. *Physical Review Letters* **113**, 251301 (2014). 1402.4119.

[72] Strüder, L. *et al.* The European Photon Imaging Camera on XMM-Newton: The pn-CCD camera. A&A **365**, L18–L26 (2001).

[73] Turner, M. J. L. *et al.* The European Photon Imaging Camera on XMM-Newton: The MOS cameras : The MOS cameras. A&A **365**, L27–L35 (2001). astro-ph/0011498.

[74] Jansen, F. *et al.* XMM-Newton observatory. I. The spacecraft and operations. A&A **365**, L1–L6 (2001).

[75] de Plaa, J. *et al.* Constraining supernova models using the hot gas in clusters of galaxies. A&A **465**, 345–355 (2007). astro-ph/0701553.

[76] Garmire, G. P., Bautz, M. W., Ford, P. G., Nousek, J. A. & Ricker, G. R., Jr. Advanced CCD imaging spectrometer (ACIS) instrument on the Chandra X-ray Observatory. In Truemper, J. E. & Tananbaum, H. D. (eds.) *X-Ray and Gamma-Ray Telescopes and Instruments for Astronomy.*, vol. 4851 of Proc. SPIE, 28–44 (2003).

[77] Weisskopf, M. C., Tananbaum, H. D., Van Speybroeck, L. P. & O'Dell, S. L. Chandra X-ray Observatory (CXO): overview. In Truemper, J. E. & Aschenbach, B. (eds.) *X-Ray Optics, Instruments, and Missions III*, vol. 4012 of Proc. SPIE, 2–16 (2000). astro-ph/0004127.

[78] Riemer-Sørensen, S. Constraints on the presence of a 3.5 keV dark matter emission line from Chandra observations of the Galactic centre. A&A **590**, A71 (2016). 1405.7943.

[79] Jeltema, T. & Profumo, S. Discovery of a 3.5 keV line in the Galactic Centre and a critical look at the origin of the line across astronomical targets. MNRAS **450**, 2143–2152 (2015). 1408.1699.

[80] Boyarsky, A., Franse, J., Iakubovskyi, D. & Ruchayskiy, O. Checking the Dark Matter Origin of a 3.53 keV Line with the Milky Way Center. *Physical Review Letters* **115**, 161301 (2015). 1408.2503.

[81] Carlson, E., Jeltema, T. & Profumo, S. Where do the 3.5 keV photons come from? A morphological study of the Galactic Center and of Perseus. JCAP **2**, 9 (2015). 1411.1758.

[82] Lovell, M. R., Bertone, G., Boyarsky, A., Jenkins, A. & Ruchayskiy, O. Decaying dark matter: the case for a deep X-ray observation of Draco. MNRAS **451**, 1573–1585 (2015). 1411.0311.

[83] Urban, O. *et al.* A Suzaku search for dark matter emission lines in the X-ray brightest galaxy clusters. MNRAS **451**, 2447–2461 (2015). 1411.0050.

[84] Franse, J. *et al.* Radial Profile of the 3.55 keV line out to R_{200} in the Perseus Cluster. *ArXiv e-prints* (2016). 1604.01759.

[85] Iakubovskyi, D., Bulbul, E., Foster, A. R., Savchenko, D. & Sadova, V. Testing the origin of ∼3.55 keV line in individual galaxy clusters

observed with XMM-Newton. *ArXiv e-prints* (2015). 1508.05186.

[86] Navarro, J. F., Frenk, C. S. & White, S. D. M. The Structure of Cold Dark Matter Halos. ApJ **462**, 563 (1996). astro-ph/9508025.

[87] Navarro, J. F., Frenk, C. S. & White, S. D. M. A Universal Density Profile from Hierarchical Clustering. ApJ **490**, 493–+ (1997). astro-ph/9611107.

[88] Smith, M. C. *et al.* The RAVE survey: constraining the local Galactic escape speed. MNRAS **379**, 755–772 (2007). astro-ph/0611671.

[89] Ruchayskiy, O. *et al.* Searching for decaying dark matter in deep XMM-Newton observation of the Draco dwarf spheroidal. MNRAS **460**, 1390–1398 (2016). 1512.07217.

[90] Bulbul, E. *et al.* Searching for the 3.5 keV Line in the Stacked Suzaku Observations of Galaxy Clusters. *ArXiv e-prints* (2016). 1605.02034.

[91] Koyama, K. *et al.* X-Ray Imaging Spectrometer (XIS) on Board Suzaku. PASJ **59**, 23–33 (2007).

[92] Mitsuda, K. *et al.* The X-Ray Observatory Suzaku. PASJ **59**, 1–7 (2007).

[93] Boyarsky, A., Franse, J., Iakubovskyi, D. & Ruchayskiy, O. Comment on the paper "Dark matter searches going bananas: the contribution of Potassium (and Chlorine) to the 3.5 keV line" by T. Jeltema and S. Profumo. *ArXiv e-prints* (2014). 1408.4388.

[94] Hofmann, F., Sanders, J. S., Nandra, K., Clerc, N. & Gaspari, M. 7.1 keV sterile neutrino constraints from X-ray observations of 33 clusters of galaxies with Chandra ACIS. *ArXiv e-prints* (2016). 1606.04091.

[95] Boyarsky, A., Ruchayskiy, O., Iakubovskyi, D., Maccio', A. V. & Malyshev, D. New evidence for dark matter. *ArXiv e-prints* (2009). 0911.1774.

[96] Iakubovskyi, D. A. New emission line at ˜3.5 keV - observational status, connection with radiatively decaying dark matter and directions for future studies. *Advances in Astronomy and Space Physics* **4**, 9–14 (2014). 1410.2852.

[97] Geringer-Sameth, A., Koushiappas, S. M. & Walker, M. Dwarf Galaxy Annihilation and Decay Emission Profiles for Dark Matter Experiments. ApJ **801**, 74 (2015). 1408.0002.

[98] Jeltema, T. E. & Profumo, S. Searching for Dark Matter with X-Ray Observations of Local Dwarf Galaxies. ApJ **686**, 1045–1055 (2008). 0805.1054.

[99] Bulbul, E. *et al.* Comment on "Dark matter searches going bananas: the contribution of Potassium (and Chlorine) to the 3.5 keV line". *ArXiv e-prints* (2014). 1409.4143.

[100] Jeltema, T. & Profumo, S. Reply to Two Comments on "Dark matter searches going bananas the contribution of Potassium (and Chlorine) to the 3.5 keV line". *ArXiv e-prints* (2014). 1411.1759.

[101] Romano, D., Karakas, A. I., Tosi, M. & Matteucci, F. Quantifying the uncertainties of chemical evolution studies. II. Stellar yields. A&A **522**, A32 (2010). 1006.5863.

[102] Phillips, K. J. H., Sylwester, B. & Sylwester, J. The X-Ray Line Feature at 3.5 KeV in Galaxy Cluster Spectra. ApJ **809**, 50 (2015). 1507.04619.

[103] Iakubovskyi, D. Checking the potassium origin of the new emission line at 3.5 keV using the K XIX line complex at 3.7 keV. MNRAS **453**, 4097–4101 (2015). 1507.02857.

[104] Gu, L. *et al.* A novel scenario for the possible X-ray line feature at ˜3.5 keV. Charge exchange with bare sulfur ions. A&A **584**, L11 (2015). 1511.06557.

[105] Cavaliere, A. & Fusco-Femiano, R. X-rays from hot plasma in clusters of galaxies. A&A **49**, 137–144 (1976).

[106] Canizares, C. R. *et al.* The Chandra High-Energy Transmission Grating: Design, Fabrication, Ground Calibration, and 5 Years in Flight. PASP **117**, 1144–1171 (2005). astro-ph/0507035.

[107] Dewey, D. Extended Source Analysis for Grating Spectrometers. In Branduardi-Raymont, G. (ed.) *High Resolution X-ray Spectroscopy with XMM-Newton and Chandra* (2002).

[108] Mitsuda, K. *et al.* Soft x-ray spectrometer (SXS): the high-resolution cryogenic spectrometer onboard ASTRO-H. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, vol. 9144 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 2 (2014).

[109] Kelley, R. L. & Mitsuda, K. Progress report on the Astro-H Soft X-Ray Spectrometer. In *AAS/High Energy Astrophysics Division*, vol. 15 of *AAS/High Energy Astrophysics Division*, 206.02 (2016).

[110] Hitomi Collaboration. The quiescent intracluster medium in the core of the Perseus cluster. *Nature* **535**, 117–121 (2016).

[111] Takahashi, T. *et al.* The ASTRO-H X-ray astronomy satellite. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, vol. 9144 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 25 (2014). 1412.1356.

[112] Figueroa-Feliciano, E. *et al.* Searching for keV Sterile Neutrino Dark Matter with X-ray Microcalorimeter Sounding Rockets. *ArXiv e-prints* (2015). 1506.05519.

[113] Barret, D. *et al.* The Hot and Energetic Universe: The X-ray Integral Field Unit (X-IFU) for Athena+. *ArXiv e-prints* (2013).

1308.6784.

[114] Ravera, L. *et al.* The X-ray Integral Field Unit (X-IFU) for Athena. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, vol. 9144 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 2 (2014).

[115] Nandra, K. *et al.* The Hot and Energetic Universe: A White Paper presenting the science theme motivating the Athena+ mission. *ArXiv e-prints* (2013). 1306.2307.

[116] Barret, D. *et al.* Athena+: The first Deep Universe X-ray Observatory. In Cambresy, L., Martins, F., Nuss, E. & Palacios, A. (eds.) *SF2A-2013: Proceedings of the Annual meeting of the French Society of Astronomy and Astrophysics*, 447–453 (2013). 1310.3814.

[117] Prokhorov, D. & Silk, J. Can the Excess in the Fe XXVI Lyγ Line from the Galactic Center Provide Evidence for 17 keV Sterile Neutrinos? ApJ **725**, L131–L134 (2010). 1001.0215.

[118] Koyama, K. *et al.* Iron and Nickel Line Diagnostics for the Galactic Center Diffuse Emission. PASJ **59**, 245–255 (2007). astro-ph/0609215.

[119] Koyama, K. *et al.* ASTRO-H White Paper - Plasma Diagnostic and Dynamics of the Galactic Center Region. *ArXiv e-prints* (2014). 1412.1170.

[120] Foster, A. *et al.* Using the new AtomDB 3.0: Non-Equilibrium Plasma Analysis. In *AAS/High Energy Astrophysics Division*, vol. 14 of *AAS/High Energy Astrophysics Division*, #115.06 (2014).

[121] Asaka, T. & Shaposhnikov, M. The @nMSM, dark matter and baryon asymmetry of the universe [rapid communication]. *Physics Letters B* **620**, 17–26 (2005). hep-ph/0505013.

[122] Asaka, T. & Shaposhnikov, M. The nuMSM, dark matter and baryon asymmetry of the universe. *Phys. Lett. B* **620**, 17–26 (2005). hep-ph/0505013.

[123] Boyarsky, A., Ruchayskiy, O. & Shaposhnikov, M. The Role of Sterile Neutrinos in Cosmology and Astrophysics. *Annual Review of Nuclear and Particle Science* **59**, 191–214 (2009). 0901.0011.

[124] Dodelson, S. & Widrow, L. M. Sterile neutrinos as dark matter. *Physical Review Letters* **72**, 17–20 (1994). hep-ph/9303287.

[125] Abazajian, K. Production and evolution of perturbations of sterile neutrino dark matter. Phys. Rev. D **73**, 063506 (2006). astro-ph/0511630.

[126] Asaka, T., Laine, M. & Shaposhnikov, M. On the hadronic contribution to sterile neutrino production. *Journal of High Energy Physics* **6**, 53 (2006). hep-ph/0605209.

[127] Viel, M., Lesgourgues, J., Haehnelt, M. G., Matarrese, S. & Riotto, A. Constraining warm dark matter candidates including sterile neutrinos and light gravitinos with wmap and the Lyman- alpha forest. *Phys. Rev.* **D71**, 063534 (2005). astro-ph/0501562.

[128] Viel, M., Lesgourgues, J., Haehnelt, M. G., Matarrese, S. & Riotto, A. Can sterile neutrinos be ruled out as warm dark matter candidates? *Phys. Rev. Lett.* **97**, 071301 (2006). astro-ph/0605706.

[129] Seljak, U., Makarov, A., McDonald, P. & Trac, H. Can Sterile Neutrinos Be the Dark Matter? *Physical Review Letters* **97**, 191303 (2006). astro-ph/0602430.

[130] Viel, M. *et al.* How cold is cold dark matter? Small scales constraints from the flux power spectrum of the high-redshift Lyman-alpha forest. *Phys. Rev. Lett.* **100**, 041304 (2008). 0709.0131.

[131] Boyarsky, A., Lesgourgues, J., Ruchayskiy, O. & Viel, M. Lyman-α constraints on warm and on warm-plus-cold dark matter models. *Journal of Cosmology and Astro-Particle Physics* **5**, 12–+ (2009). 0812.0010.

[132] Boyarsky, A., Lesgourgues, J., Ruchayskiy, O. & Viel, M. Realistic Sterile Neutrino Dark Matter with KeV Mass does not Contradict Cosmological Bounds. *Physical Review Letters* **102**, 201304–+ (2009). 0812.3256.

[133] Boyarsky, A., Ruchayskiy, O. & Iakubovskyi, D. A lower bound on the mass of dark matter particles. *Journal of Cosmology and Astro-Particle Physics* **3**, 5–+ (2009). 0808.3902.

[134] Gorbunov, D., Khmelnitsky, A. & Rubakov, V. Constraining sterile neutrino dark matter by phase-space density observations. *JCAP* **0810**, 041 (2008). 0808.3910.

[135] Angus, G. W. A lower limit on the dark particle mass from dSphs. JCAP **3**, 26 (2010). 0907.1526.

[136] Shao, S., Gao, L., Theuns, T. & Frenk, C. S. The phase-space density of fermionic dark matter haloes. MNRAS **430**, 2346–2357 (2013). 1209.5563.

[137] Shi, X. & Fuller, G. M. New Dark Matter Candidate: Nonthermal Sterile Neutrinos. *Physical Review Letters* **82**, 2832–2835 (1999). astro-ph/9810076.

[138] Laine, M. & Shaposhnikov, M. Sterile neutrino dark matter as a consequence of nuMSM-induced lepton asymmetry. *Journal of Cosmology and Astro-Particle Physics* **6**, 31–+ (2008). 0804.4543.

[139] Abazajian, K. N. Resonantly Produced 7 keV Sterile Neutrino Dark Matter Models and the Properties of Milky Way Satellites. *Physical Review Letters* **112**, 161303 (2014). 1403.0954.

[140] Ghiglieri, J. & Laine, M. Improved determina-

tion of sterile neutrino dark matter spectrum. *Journal of High Energy Physics* **11**, 171 (2015). 1506.06752.

[141] Venumadhav, T., Cyr-Racine, F.-Y., Abazajian, K. N. & Hirata, C. M. Sterile neutrino dark matter: A tale of weak interactions in the strong coupling epoch. *ArXiv e-prints* (2015). 1507.06655.

[142] Liao, W. keV scale ν_R dark matter and its detection in β decay experiments. Phys. Rev. D **82**, 073001 (2010). 1005.3351.

[143] Liao, W., Wu, X.-H. & Zhou, H. Electron events from the scattering with solar neutrinos in the search of keV scale sterile neutrino dark matter. Phys. Rev. D **89**, 093017 (2014). 1311.6075.

[144] Mertens, S. *et al.* Wavelet approach to search for sterile neutrinos in tritium β -decay spectra. Phys. Rev. D **91**, 042005 (2015). 1410.7684.

[145] Dragoun, O. & Vénos, D. Searches for Active and Sterile Neutrinos in Beta-Ray Spectra. *ArXiv e-prints* (2015). 1504.07496.

[146] Adhikari, R. *et al.* A White Paper on keV Sterile Neutrino Dark Matter. *ArXiv e-prints* (2016). 1602.04816.

[147] Bonivento, W. *et al.* Proposal to Search for Heavy Neutral Leptons at the SPS. *ArXiv e-prints* (2013). 1310.1762.

[148] Alekhin, S. *et al.* A facility to Search for Hidden Particles at the CERN SPS: the SHiP physics case. *ArXiv e-prints* (2015). 1504.04855.

[149] Blondel, A., Graverini, E., Serra, N., Shaposhnikov, M. & for the FCC-ee study team. Search for Heavy Right Handed Neutrinos at the FCC-ee. *ArXiv e-prints* (2014). 1411.5230.

[150] Macciò, A. V., Ruchayskiy, O., Boyarsky, A. & Muñoz-Cuartas, J. C. The inner structure of haloes in cold+warm dark matter models. MNRAS **428**, 882–890 (2013). 1202.2858.

[151] Lovell, M. R. *et al.* The properties of warm dark matter haloes. MNRAS **439**, 300–317 (2014). 1308.1399.

[152] Viel, M., Becker, G. D., Bolton, J. S. & Haehnelt, M. G. Warm dark matter as a solution to the small scale crisis: New constraints from high redshift Lyman-α forest data. Phys. Rev. D **88**, 043502 (2013). 1306.2314.

[153] Schneider, A. Structure formation with suppressed small-scale perturbations. MNRAS **451**, 3117–3130 (2015). 1412.2133.

[154] Merle, A. & Schneider, A. Production of Sterile Neutrino dark matter and the 3.5 keV line. *Physics Letters B* **749**, 283–288 (2015). 1409.6311.

[155] Bozek, B. *et al.* Resonant sterile neutrino dark matter in the local and high-z Universe. MNRAS **459**, 1489–1504 (2016). 1512.04544.

[156] Lovell, M. R. *et al.* Satellite galaxies in semi-analytic models of galaxy formation with sterile neutrino dark matter. MNRAS (2016). 1511.04078.

[157] Bose, S. *et al.* The Copernicus Complexio: statistical properties of warm dark matter haloes. MNRAS **455**, 318–333 (2016). 1507.01998.

[158] Wang, M.-Y., Strigari, L. E., Lovell, M. R., Frenk, C. S. & Zentner, A. R. Mass assembly history and infall time of the Fornax dwarf spheroidal galaxy. MNRAS **457**, 4248–4261 (2016). 1509.04308.

[159] Li, R. *et al.* Constraints on the identity of the dark matter from strong gravitational lenses. MNRAS (2016). 1512.06507.

[160] Bose, S. *et al.* Substructure and galaxy formation in the Copernicus Complexio warm dark matter simulations. *ArXiv e-prints* (2016). 1604.07409.

[161] Ludlow, A. D. *et al.* The mass-concentration-redshift relation of cold and warm dark matter haloes. MNRAS **460**, 1214–1232 (2016). 1601.02624.

[162] Schneider, A. Astrophysical constraints on resonantly produced sterile neutrino dark matter. JCAP **4**, 059 (2016). 1601.07553.

[163] Kamada, A., Inoue, K. T. & Takahashi, T. Constraints on mixed dark matter from anomalous strong lens systems. *ArXiv e-prints* (2016). 1604.01489.

[164] Rudakovskyi, A. & Iakubovskyi, D. Influence of ˜7 keV sterile neutrino dark matter on the process of reionization. JCAP **6**, 017 (2016). 1604.01341.

[165] Bose, S., Frenk, C. S., Jun, H., Lacey, C. G. & Lovell, M. R. Reionisation in sterile neutrino cosmologies. *ArXiv e-prints* (2016). 1605.03179.

[166] Finkbeiner, D. P. & Weiner, N. An X-Ray Line from eXciting Dark Matter. *ArXiv e-prints* (2014). 1402.6671.

[167] Cline, J. M., Liu, Z., Moore, G. D., Farzan, Y. & Xue, W. 3.5 keV x rays as the "21 cm line" of dark atoms, and a link to light sterile neutrinos. Phys. Rev. D **89**, 121302 (2014). 1404.3729.

[168] Okada, H. & Toma, T. 3.55 keV X-ray line signal from excited dark matter in radiative neutrino model. *Physics Letters B* **737**, 162–166 (2014). 1404.4795.

[169] Cline, J. M. & Frey, A. R. Nonabelian dark matter models for 3.5 keV X-rays. JCAP **10**, 13 (2014). 1408.0233.

[170] Boddy, K. K., Feng, J. L., Kaplinghat, M., Shadmi, Y. & Tait, T. M. P. Strongly interacting dark matter: Self-interactions and keV lines. Phys. Rev. D **90**, 095016 (2014). 1408.6532.

[171] Schutz, K. & Slatyer, T. R. Self-scattering for

Dark Matter with an excited state. JCAP **1**, 21 (2015). 1409.2867.

[172] Cline, J. M. & Frey, A. R. Consistency of dark matter interpretations of the 3.5 keV x-ray line. Phys. Rev. D **90**, 123537 (2014). 1410.7766.

[173] Berlin, A., DiFranzo, A. & Hooper, D. 3.55 keV line from exciting dark matter without a hidden sector. Phys. Rev. D **91**, 075018 (2015). 1501.03496.

[174] D'Eramo, F., Hambleton, K., Profumo, S. & Stefaniak, T. Dark matter inelastic upscattering with the interstellar plasma: A new source of x-ray lines, including at 3.5 keV. Phys. Rev. D **93**, 103011 (2016). 1603.04859.

[175] Dudas, E., Heurtier, L. & Mambrini, Y. Generating x-ray lines from annihilating dark matter. Phys. Rev. D **90**, 035002 (2014). 1404.1927.

[176] Frandsen, M. T., Sannino, F., Shoemaker, I. M. & Svendsen, O. X-ray lines from dark matter: the good, the bad, and the unlikely. JCAP **5**, 33 (2014). 1403.1570.

[177] Baek, S., Ko, P. & Park, W.-I. The 3.5 keV X-ray line signature from annihilating and decaying dark matter in Weinberg model. *ArXiv e-prints* (2014). 1405.3730.

[178] Mambrini, Y. & Toma, T. X-ray lines and self-interacting dark matter. *European Physical Journal C* **75**, 570 (2015). 1506.02032.

[179] Cicoli, M., Conlon, J. P., Marsh, M. C. D. & Rummel, M. 3.55 keV photon line and its morphology from a 3.55 keV axionlike particle line. Phys. Rev. D **90**, 023540 (2014). 1403.2370.

[180] Conlon, J. P. & Day, F. V. 3.55 keV photon lines from axion to photon conversion in the Milky Way and M31. JCAP **11**, 33 (2014). 1404.7741.

[181] Conlon, J. P. & Powell, A. J. A 3.55 keV line from DM $\rightarrow a + \gamma$: predictions for cool-core and non-cool-core clusters. JCAP **1**, 19 (2015). 1406.5518.

[182] Alvarez, P. D., Conlon, J. P., Day, F. V., Marsh, M. C. D. & Rummel, M. Observational consistency and future predictions for a 3.5 keV ALP to photon line. JCAP **4**, 13 (2015). 1410.1867.

[183] Berg, M. *et al.* Searches for Axion-Like Particles with NGC1275: Observation of Spectral Modulations. *ArXiv e-prints* (2016). 1605.01043.

[184] Kitayama, T. *et al.* ASTRO-H White Paper - Clusters of Galaxies and Related Science. *ArXiv e-prints* (2014). 1412.1176.

[185] Speckhard, E. G., Ng, K. C. Y., Beacom, J. F. & Laha, R. Dark Matter Velocity Spectroscopy. *Physical Review Letters* **116**, 031301 (2016). 1507.04744.

[186] Merloni, A. *et al.* eROSITA Science Book: Mapping the Structure of the Energetic Universe. *ArXiv e-prints* (2012). 1209.3114.

[187] Zane, S. *et al.* The large area detector of LOFT: the Large Observatory for X-ray Timing. In *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, vol. 9144 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 2 (2014). 1408.6539.

[188] Zandanel, F., Weniger, C. & Ando, S. The role of the eROSITA all-sky survey in searches for sterile neutrino dark matter. JCAP **9**, 060 (2015). 1505.07829.

[189] Boyarsky, A., Iakubovskyi, D. & Ruchayskiy, O. Next decade of sterile neutrino studies. *Physics of the Dark Universe* **1**, 136–154 (2012). 1306.4954.

[190] Neronov, A. & Malyshev, D. Toward a full test of the ν MSM sterile neutrino dark matter model with Athena. Phys. Rev. D **93**, 063518 (2016). 1509.02758.